

## First record of the freshwater sawfish, *Pristis microdon*, from southwestern Australian waters

Justin A. Chidlow

Department of Fisheries, Western Australian Marine Research Laboratories,  
PO Box 20, North Beach, Western Australia 6920  
Email: jchidlow@fish.wa.gov.au

Sawfishes (family Pristidae) are large (up to 7m) modified batoids with a blade-like snout edged with pairs of rostral teeth. They occur worldwide in sub-tropical and tropical shallow coastal sea, estuaries and freshwater systems (Last and Stevens 1994; Compagno and Last 1998). There are between five and seven recognised species worldwide, with five species represented in Australian waters (Last and Stevens 1994). Sawfish populations have been extirpated from many parts of their original global range by gillnetting and trawling and are easily entangled in nets by their toothed rostra (Simpfendorfer 2000). The little that is known about the biology of sawfish suggests they have low rates of reproduction (Tanaka 1991; Compagno and Last 1998; Wilson 1999; Simpfendorfer 2000; Thorburn *et al.* 2004). This combined with their susceptibility to fishing gear, make sawfish a high risk species and all have subsequently been listed globally as critically endangered under the IUCN Red List Assessment 2006 (Compagno *et al.* 2006).

### *Pristis microdon* Latham, 1794

*Pristis microdon* is a medium to large sawfish that in Australia grows to at least 361cm TL (Tanaka 1991), but is reported to reach up to 700cm TL in other locations (Last and Stevens 1994). They are born at around 50cm in length after a five month gestation period, with litter sizes ranging between 1 and 12 (Wilson 1999). In the western Atlantic *P. microdon* matures at between 240cm and 300cm TL (Compagno and Last 1998). Tanaka (1991) reported two male specimens from New Guinea, one measuring 247cm that was immature and a 361cm specimen that was mature. In Australian waters, *P. microdon* feeds on fish such as catfish, small crustaceans and molluscs (Allen 1982; Cliff and Wilson 1994; Pogonoski *et al.* 2002; Thorburn *et al.* 2004).

*Pristis microdon* occurs inshore and in intertidal areas and is usually found in freshwater drainages, lakes and estuaries where it can penetrate as far as 400km from the coast (Morgan *et al.* 2004). In the Indo-West Pacific it ranges from New Guinea, SE Asia, northern Australia and west to South Africa

(Last and Stevens 1994; Compagno and Last 1998). *Pristis microdon* may also occur in the Atlantic and eastern Pacific if *P. perotteti* Müller & Henle, 1841 and *P. zephyreus* Jordan & Starks in Jordan, 1895 are synonymised with this species (Compagno and Last 1998). In Australia, the freshwater sawfish is known to occur in the Ord, Durack and Fitzroy Rivers (Western Australia), the Adelaide, Victoria and Daly Rivers (Northern Territory), and the Gilbert, Mitchell, Norman and Leichhardt Rivers (Queensland) (Last and Stevens 1994; Pogonoski *et al.* 2002; Thorburn *et al.* 2004). Only recently has *P. microdon* been reported from marine waters (Thorburn *et al.* 2004).

### Southwestern Australian *P. microdon*

A female *P. microdon* was captured by a commercial shark fisher operating demersal gillnets in southwestern Australian waters on the 5<sup>th</sup> of February 2003. The capture location was approximately six miles east of Cape Naturaliste (33°31'S, 115°07'E) in 32m of water. The sawfish was estimated to be 3.5m in length TL when landed and appeared to be healthy. The specimen was processed and the fisher retained the remaining trunk, fins and saw. I positively identified the processed sawfish as *P. microdon* using an identification key provided by Last and Stevens (1994).

The partial length (origin of the first dorsal fin to the insertion point of the second dorsal fin) was 95cm (approximate as the trunk had been cut in half). The rostral saw length was 79cm with 19 pairs of teeth that extended to the basal quarter of the saw (Figure 1). The interspace between rostral teeth at the base of the saw was 4cm, and 3cm between the teeth at the tip of the saw (Figure 1). A groove was present along the posterior margin of all rostral teeth. The origin of the first dorsal-fin was located anterior to the pelvic-fin origin and the height of the first dorsal-fin was 32cm. The second dorsal-fin height was 31cm. The ventral lobe of the caudal-fin was small, but distinct. The upper and lower postventral caudal-fin margins measured 44.5cm and 11.5cm respectively.




Figure 1 Rostral saw from a female *Pristis microdon*, measuring approximately 350cm in total length, captured off Cape Naturaliste, Western Australia. See text for description of measurements.

This record of *P. microdon* from southwestern Australia extends the range of the species approximately 1600 km south of its previously known southern limit, Cape Keraudren, Western Australia (Thorburn *et al.* 2004) and provides further confirmation that *P. microdon* utilizes marine waters.

#### ACKNOWLEDGEMENTS

I wish to thank J. Nelson who thoughtfully informed the Shark Research Section, Department of Fisheries WA of the capture, and provided assistance in identifying and collecting data from the specimen. I would also like to thank P. Last from CSIRO Marine Research, Hobart, R. McAuley from Department of Fisheries WA, Perth and C. Simpfendorfer from the Mote Marine Laboratory, Florida for their assistance in positively identifying the specimen.

#### REFERENCES

- Allen, G.R. (1982). *A field guide to inland fishes of Western Australia*. Western Australian Museum, Perth, Western Australia. 86pp.
- Cliff, G. and Wilson, G. (1994). Natal sharks board's guide to sharks and other marine animals. *Natal Sharks Board*, p33.
- Compagno, L.J.V. and Last, J.D. (1999). Pristiformes: Pristidae. In K.E. Carpenter and V.H. Niem (eds). *FAO species identification guide for fishery purposes. The living marine resources of the Western Central Pacific. Volume 3. Batoid fishes, chimaeras and bony fishes part 1 (Elopidae to Linophrynidae)*. Pp. 1410–1417. FAO, Rome.
- Compagno, L.J.V., Cook, S.F. and Fowler, S.L. (2006). *Pristis microdon*. In IUCN 2006 IUCN Red List of Threatened Species.
- Jordan, D.S. and Starks, E.C. (1895). In *The fishes of Sinaloa*. D.S. Jordan. Proceedings of the California Academy of Sciences (Ser. 2) 377–514.
- Last, P.R. and Stevens, J.D. (1994). *Sharks and rays of Australia*. CSIRO, Melbourne, 513 pp.
- Latham, J. (1794). An essay on the various species of sawfish. *Transactions of the Linnean Society of London* 2 (25): 273–282.
- Morgan, D.L., Allen, M.G., Bedford, P. and Horstman, M. (2004). Fish fauna of the Fitzroy River in the Kimberley region of Western Australia – including Bunuba, Gooniyandi, Ngarinyin, Nyikina and Walmajarri Aboriginal names. *Records of the Western Australian Museum* 22:147–161.
- Müller, J. and Henle, F.G.J. (1841). Systematische Beschreibung der Plagiostomen Berlin. Plagiostomen i–xxii + 1–200.
- Pogonoski, J.J., Pollard, D.A. and Paxton, J.R. (2002). Conservation overview and action plan for Australian threatened and potentially threatened marine and estuarine fishes. *Environment Australia*, February 2002.
- Simpfendorfer, C.A. (2000). Predicting population recovery rates for endangered western Atlantic sawfishes using demographic analysis. *Environmental Biology of Fishes* 58: 371–377.
- Tanaka, S. (1991). Age estimation of freshwater sawfish and sharks in northern Australia and Papua New Guinea. The University Museum, University of Tokyo. *Nature and Culture* 3: 71–82.
- Thorburn, D., Morgan, D., Gill, H., Johnson, M., Wallace-Smith, H., Vigilante, T., Gorring, A., Croft, I. and Fenton, J. (2004). Biology and cultural significance of the freshwater sawfish (*Pristis microdon*) in the Fitzroy River, Kimberley, Western Australia. *Report to the Threatened Species Network* 2004. 57 pp.
- Thorson, T.B. (1982). The impact of commercial exploitation on sawfish and shark populations in Lake Nicaragua. *Fisheries* 7(2): 2–10.
- Wilson, D. (1999). Freshwater sawfish *Pristis microdon*. Australia New Guinea Fishes Associations' A-Z notebook of native freshwater fish. *ANGFA Bulletin* 41.